arXiv:1612.05821v1 [math.FA] 17 Dec 2016

Invariant subspaces for commuting operators in a real Banach space

Victor Lomonosov and Victor Shulman

20 декабря 2016 г.

Abstract.

It is proved that a commutative algebra A of operators in a reflexive real Banach space has an invariant subspace if each operator $T \in A$ satisfies the condition

$$\|1 - \varepsilon T^2\|_e \leq 1 + o(\varepsilon) \text{ when } \varepsilon \searrow 0,$$

where $\|\cdot\|_e$ is the essential norm. This implies the existence of an invariant subspace for every commutative family of essentially selfadjoint operators in a real Hilbert space.

1 Introduction

One of the most known unsolved problems in the invariant subspace theory is the question of existence of a (non-trivial, closed) invariant subspace for an operator T with compact imaginary part (= essentially selfadjoint operator = compact perturbation of a selfadjoint operator). There is a lot of papers concerning this subject; we only mention that the answer is affirmative for perturbations by operators from Shatten - von Neumann class $\mathfrak{S}_p$ (Livshits [1] for $p = 1$, Sahnovich [2] for $p = 2$, Gohberg and Krein [3], Macaev [4], Schwartz [5] — for arbitrary p), or, more generally, from the Macaev ideal (Macaev [6]). But the general question is still open.

It was proved in [7] that an essentially self-adjoint operator in a complex Hilbert space has an invariant real subspace. Then in [8] the following general theorem of Burnside type was proved:

Theorem 1.1 *Suppose that an algebra A of bounded operators in a (real or complex) Banach space X is not dense in the algebra $\mathcal{B}(X)$ of all bounded operators on X with respect to the weak operator topology (WOT). Then there are non-zero $x \in X^{**}, y \in X^*$, such that*

$$|(x, T^*y)| \leq \|T\|_e \text{ for all } T \in A, \tag{1.1}$$

where $\|T\|_e$ is the essential norm of T, that is $\|T\|_e = \inf\{\|T + K\| : K \in \mathcal{K}(X)\}$. Here $\mathcal{K}(X)$ is the ideal of all compact operators in X.

Using this result and developing a special variational techniques, Simonic [9] has obtained a significant progress in the topic: he proved that each essentially selfadjoint operator in a real Hilbert space has invariant subspace. Deep results based on Theorem 1.1 were proved then by Atzmon

[10], Atzmon, Godefroy and Kalton [11], Grivaux [12] and other mathematicians. Here we show that every commutative family of essentially selfadjoint operators in a real Hilbert space has an invariant subspace, and consider some analogs of this result for operators in Banach spaces. Our proof is very simple and short but it heavily depends on Theorem 1.1. More precisely, we use the following improvement of Theorem 1.1 obtained in [13]:

Theorem 1.2 *If an algebra $A \subset \mathcal{B}(X)$ is not WOT-dense in $\mathcal{B}(X)$, then there are non-zero vectors $x \in X^{**}, y \in X^*$, such that $(x, y) \geq 0$ and*

$$|(x, T^*y)| \leq \|T\|_e(x, y), \text{ for all } T \in A. \tag{1.2}$$

Let us mention that if A contains a non-zero compact operator then by [14] A has a non-trivial invariant subspace L in X (so x can be chosen from L and y can be chosen from $L^\perp$).

The original proof of Theorem 1.1 was essentially simplified by Lindstrom and Shluchtermann [15]. For completeness, we give at the end of the paper a short proof of Theorem 1.2, unifying arguments from [15] and [13] (and correcting some inaccuracy in [13]) — in this form it was not published before.

2 Main results

In this section X is a real Banach space (complex spaces are considered as real ones). The standard epimorphism from $\mathcal{B}(X)$ to the Calkin algebra $\mathcal{C}(X) = \mathcal{B}(X)/\mathcal{K}(X)$ is denoted by π.

Let us say that an element a of a unital normed algebra is *positive*, if $\|1 - \varepsilon a\| \leq 1 + o(\varepsilon)$ for $\varepsilon > 0$, $\varepsilon \to 0$. And let us say that an element a is *real*, if a^2 is positive. An operator $T \in \mathcal{B}(X)$ is *essentially real*, if $\pi(T)$ is a real element of $\mathcal{C}(X)$.

Clearly all selfadjoint operators in a Hilbert space are real. It is not difficult to check that Hermitian operators in a complex Banach space (defined by the condition $\|\exp(itT)\| = 1$ for $t \in \mathbb{R}$) are real. Many other operators, for instance, all involutions and all nilpotents of index two are also real. So we can see that the class of essentially real operators is very wide.

Theorem 2.1 *If $A \subset \mathcal{B}(X)$ is a commutative algebra of essentially real operators, then there exists a non-trivial closed subspace of X^*, invariant for the algebra $A^* = \{T^* : T \in A\}$.*

Proof. Note that the set of all positive elements of a Banach algebra is a convex cone. Moreover this cone is norm-closed. Indeed let $a = \lim_{n\to\infty} a_n$ where all a_n are positive. If a is not positive then there is a sequence $\varepsilon_n \to 0$ and a number $C > 0$, such that $\|1 - \varepsilon_n a\| > 1 + C\varepsilon_n$ for all n. Taking k with $\|a - a_k\| < C/2$, we get that $\|1 - \varepsilon_n a_k\| > 1 + C\varepsilon_n - \|a - a_k\|\varepsilon_n > 1 + (C/2)\varepsilon_n$, which is a contradiction to positivity of a_k.

Hence the set of real elements is also norm closed. Applying this to $\mathcal{C}(X)$ we see that the set of essentially real operators is norm-closed as well. This allows us to assume that the algebra A is norm-closed. Obviously we may assume also that A is unital. Therefore $\exp(T) \in A$, for each $T \in A$.

Since A is commutative it is not WOT-dense in $\mathcal{B}(X)$. Applying Theorem 1.2, we find non-zero vectors $x \in X^{**}, y \in X^*$, such that the condition (1.2) holds.

Therefore, for $T \in A$ and $\varepsilon \searrow 0$, we have

$$(x, (1 - \varepsilon(T^2)^*)y) \leq \|1 - \varepsilon T^2\|_e(x, y) \leq (1 + o(\varepsilon))(x, y),$$

hence $-\varepsilon(x,(T^2)^*y) \le o(\varepsilon)(x,y)$ and $(x,(T^2)^*y) \ge 0$, because $(x,y) \ge 0$.

Since $\exp(T) = (\exp(T/2))^2$, we get that

$$(x, \exp(T^*)y) \ge 0 \text{ for } T \in A. \tag{2.1}$$

Let K be the closed convex envelope of the set $M = \{\exp(T^*)y : T \in A\} \subset X^*$. Since M is invariant under all operators $\exp(T^*)$, the same is true for K.

Since $K \ne X^*$ (by (2.1)) and K is not a singleton (otherwise we trivially have an invariant subspace $\mathbb{C}y$), the boundary ∂K of K is not a singleton. By the Bishop-Phelps theorem [17], the set of support points is dense in ∂K, so there is a non-zero functional $x_0 \in X^{**}$ supporting K in some point $0 \ne y_0 \in K$. That is

$$(x_0, y_0) \le (x_0, z) \text{ for all } z \in K.$$

By the above arguments, $\exp(T^*)y_0 \in K$ for $T \in A$, therefore

$$(x_0, y_0) \le (x_0, \exp(T^*)y_0) \text{ for all } T \in A.$$

It follows that for each $T \in A$, the function $\phi(t) = (x_0, \exp(tT^*)y_0)$ has a minimum at $t = 0$. For this reason $\phi'(0) = 0$ and $(x_0, T^*y_0) = 0$. Hence the subspace A^*y_0 is not dense in H, and its norm-closure $\overline{A^*y}$ is a non-trivial invariant subspace. ■

Corollary 2.2 *A commutative algebra of essentially real operators in a reflexive real Banach space has a non-trivial invariant subspace.*

Since the algebra generated by a commutative family of essentially selfadjoint operators consists of essentially selfadjoint operators we get the following result:

Corollary 2.3 *Any commutative family of essentially selfadjoint operators in a real Hilbert space has a non-trivial invariant subspace.*

Returning to individual criteria of continuity let us consider the class (E) of operators T such that all polynomials $p(T)$ of T are essentially real.

Corollary 2.4 *Each operator $T \in (E)$ in a reflexive real Banach space has a non-trivial invariant subspace.*

Atzmon, Godefroy and Kalton [11] introduced the class (S) of all operators, satisfying the condition

$$\|p(T)\|_e \le \sup\{|p(t)| : t \in L\}, \text{ for each polynomial } p, \tag{2.2}$$

where L is a compact subset of $\mathbb{R}$. It was proved in [11] that all operators in (S) have invariant subspaces. It is not difficult to see that $(S) \subset (E)$ (indeed if $T \in (S)$ then $\|1 - \varepsilon p(T)^2\|_e \le \sup\{|1 - \varepsilon p(t)^2| : t \in L\} \le 1$ if ε is sufficiently small) , so this result follows from Corollary 2.4.

3 Proof of Theorem 1.2

Without loss of generality one can assume that A is norm-closed. Since the algebra A is not WOT-dense in $\mathcal{B}(X)$, the algebra A^* is not WOT-dense in $\mathcal{B}(X^*)$. Suppose, aiming at the contrary, that A^* is transitive (has no invariant subspaces). Set $F = \{T^* \in A^* : \|T\|_e < 1\}$ and fix $\varepsilon \in (0, \frac{1}{10})$. Choose $y_0 \in X^*$ with $\|y_0\| = 3$ and let S be the ball $\{y \in X^* : \|y - y_0\| \leq 2\}$.

Let us suppose firstly that Fy is dense in X^* for each non-zero $y \in X^*$. Then the same is true for εFy. It follows that for every $y \in S$ there exists $T_y^* \in \varepsilon F$ with $\|T_y^* y - y_0\| < 1$. By the definition of F, $T_y^* = R_y^* + K_y^*$, where $\|R_y\| < \varepsilon$, $K_y \in \mathcal{K}(X)$. Thus $\|K_y^* y - y_0\| \leq \|T_y^* y - y_0\| + \|R_y^* y\| < 1 + 5\varepsilon$ (since $\|y\| \leq 5$, for each $y \in S$). Let τ denote the (relative) *-weak topology of S, then compactness of K_y implies that K_y^* continuously maps (S, τ) to $(X^*, \|\cdot\|)$. Therefore, for each $y \in S$, there is a τ-neighborhood V_y of y such that $\|K_y^* z - y_0\| < 1 + 5\varepsilon$, for each $z \in V_y$, and $\|T_y^* z - y_0\| < 1 + 5\varepsilon + 5\varepsilon < 2$. In other words T_y^* maps V_y to S.

The sets V_y, $y \in S$, form an open covering of S. Since S is τ-compact there is a finite subcovering $\{V_{y_i} : 1 \leq i \leq n\}$. Let $\{\varphi_i : 1 \leq i \leq n\}$ be a partition of unity related to this subcovering. We define a τ-continuous map $\Phi : S \to S$ by $\Phi(y) = \sum_{i=1}^n \varphi_i(y) T_{y_i}^*(y)$. By Tichonov's Theorem, Φ has a fixed point $z \in S$. This means that $T^* z = z$, where $T^* = \sum_{i=1}^n \varphi_i(z) T_{y_i}^*$. Since the set εF is convex we get that $T^* \in \varepsilon F$ and $\|T^*\|_e \leq \|T\|_e < \frac{1}{10}$.

For this reason 1 is an eigenvalue of T^* exceeding $\|T^*\|_e$ and hence it is an isolated point in $\sigma(T^*)$. The corresponding Riesz projection is of finite rank and belongs to A^*. But it is well known (see e.g. [16, Theorem 8.2]), that a transitive algebra containing a non-zero finite rank operator is WOT-dense in the algebra of all operators. Since A^* is not WOT-dense in $\mathcal{B}(X^*)$, we obtain a contradiction.

Hence there exists $y_0 \in X^*$ such that Fy_0 is not norm-dense in X^*. Let V be the norm-closure of Fy_0. If $V = \{0\}$ then we have the inequality (1.2) with $y = y_0$ and any non-zero x.

If $V \neq \{0\}$ then V is a norm-closed convex proper subset of X^* containing more then one point. By the Bishop - Phelps Theorem [17], there are $0 \neq y \in V$ and $0 \neq x \in X^{**}$ such that $Re(x, y) = \sup\{Re(x, w) : w \in V\}$. Since the set V is invariant under multiplication by any number t with $|t| \leq 1$, we have $(x, y) \geq 0$ and $(x, y) \geq |(x, w)|$ for all $w \in V$. Since $F^2 \subset F$, we have that $Fy \subset V$ and $|(x, T^* y)| \leq (x, y)$, for any $T^* \in F$. Therefore the inequality $|(x, T^* y)| \leq \|T\|_e (x, y)$ holds for all $T \in A$. ■

Список литературы

[1] Livsic M.S., On the spectral resolution of linear non-selfadjoint operators, Mathem. Sbornik, 34(76), (1954), 145-198 (Amer. Math. Soc. Transl. (2) 5 (1957), 67-114).

[2] Sahnovich L.A., On reduction of a non-selfadjoint operators to the triangular form, Izvestia Vuzov. Mathematika, 1959,4,141-149.

[3] Gohberg I.Ts., Krein M.G., On completely continuous operators with spectrum concentrated in zero, Soviet Math. Dokl., 128 (1959),2, 227-230.

[4] Macaev V.I., On Volterra operators which are perturbations of selfadjoint ones, Soviet Math. Dokl., 139(1961),4, 810-814.

[5] Schwartz J., Subdiagonalization of operators in Hilbert space with compact imaginary part, Comm. Pure Appl. Math. 15(1962), 159-172.

[6] Macaev V.I., On a class of completely continuous operators, Soviet Math. Dokl., 139(1961),3, 548-552.

[7] Lomonosov V., On real invariant subspaces of bounded operators with compact imaginary part, Proc. Amer. Math. Soc. 115(1992), 775-777.

[8] Lomonosov V., An extension of Burnside's theorem to infinite-dimensional spaces, Israel J. Math. 75(1991), 329-339.

[9] Simonic A., An extension of Lomonosov's techniques to non-compact operators, Trans. Amer. Math. Soc. 348(1996), 975-995.

[10] Atzmon A., Reducible representations of abelian groups, Ann.Inst.Fourier 51(2001), 1407-1418.

[11] Atzmon A., Godefroy G., Kalton N.J., Invariant subspaces and the exponential map, Positivity 8(2004), no 2, 101-107.

[12] Grivaux S., Invariant subspaces for tridiagonal operators, Bull. sci. math. 126(2002), 681-691.

[13] Lomonosov V.I., Positive functionals on general operator algebras, J.Math.Anal.Appl., 245(2000), 221-224.

[14] Lomonosov V.I., Invariant subspaces for operators commuting with compact operators, Functional Analysis and Applications, 7(1973), 213-214.

[15] Lindstrom M., Shluchtermann G., Lomonosov's technique and Burnside's theorem, Canad. Math. Bull. 43(2000), 87-89.

[16] Radjavi H., Rosenthal P., "Invariant subspaces" , Springer Verlag, 1970.

[17] Bishop E., Phelps R.R., The support functionals of a convex set, Proc. Symp. of Pure Math. VII, Covexity, Amer. Math. Soc., 1963, 27-36.

Dept of Math.
Kent State University
Kent OH 44242, USA
lomonoso@math.kent.edu

Dept of Math.
Vologda State University
Vologda 160000, Russia
shulman.victor80@gmail.com